

L'exemplaire filmé fut reproduit grâce à la générosité de:




Engineering Sciences Library,
Queen's University

Les images suivantes ont été reproduites avec le plus grand soin, compte tenu de la condition et de la netteté de l'exemplaire filmé, et en conformité avec les conditions du contrat de filmage.

Les exemplaires originaux dont la couverture en papier est imprimée sont filmés en commençant par le premier plat et en terminant soit par la dernière page qui comporte une empreinte d'impression ou d'illustration, soit par le second plat, selon le cas. Tous les autres exemplaires originaux sont filmés en commençant par la première page qui comporte une empreinte d'impression ou d'illustration et en terminant par la dernière page qui comporte une telle empreinte.

Un des symboles suivants apparaîtra sur la dernière image de chaque microfiche, selon le cas: le symbole \rightarrow signifie "A SUIVRE", le symbole ∇ signifie "FIN".

Les cartes, planches, tableaux, etc., peuvent être filmés à des taux de réduction différents. Lorsque le document est trop grand pour être reproduit en un seul cliché, il est filmé à partir de l'angle supérieur gauche, de gauche à droite, et de haut en bas, en prenant le nombre d'images nécessaire. Les diagrammes suivants illustrent la méthode.



**UNIVERSITY OF TORONTO
STUDIES**

**PAPERS FROM THE
PHYSICAL LABORATORIES**

**No. 61: NEW LINES IN THE EXTREME ULTRA-VIOLET OF
CERTAIN METALS, BY D. S. AINSLIE and D. S. FULLER**

(REPRINTED FROM TRANSACTIONS OF THE ROYAL SOCIETY OF CANADA, SERIES III, Vol. XII)

**THE UNIVERSITY LIBRARY: PUBLISHED BY
THE LIBRARIAN, 1918**

UNIVERSITY OF TORONTO STUDIES
COMMITTEE OF MANAGEMENT

Chairman: SIR ROBERT ALEXANDER FALCONER, LL.D., K.C.M.G.
President of the University

PROFESSOR W. J. ALEXANDER, Ph.D.

PROFESSOR W. H. ELLIS, M.A., M.B.

PROFESSOR J. J. MACKENZIE, B.A., M.B.

PROFESSOR J. P. McMURRICH, Ph.D.

PROFESSOR G. H. NEEDLER, Ph.D.

PROFESSOR GEORGE M. WRONG, M.A.

General Editor: H. H. LANGTON, M.A.
Librarian of the University

New Lines in the Extreme Ultra-Violet of Certain Metals.

By D. S. AINSLIE, M.A., and D. S. FULLER, M.A.

University of Toronto.

Presented by PROFESSOR E. F. BURTON, Ph.D., F.R.S.C.

(Read May Meeting, 1918.)

The experiments described below form an extension of the recent work of Lyman,¹ Handke,² Wolff,³ and Saunders⁴, on the extreme ultra-violet region.

By using a fluorite vacuum spectroscope and a vacuum arc lamp, photographs of spectral lines were obtained free from the disturbance due to the absorption of the light by glass, quartz or air. The arc could be manipulated from the outside of the case of the spectrograph; the current ranged from four to fifteen amperes, according to the metal used.

For the hard metals with high melting points the terminals of the arc were made of the metal; when a soft metal was used, the terminals were made of iron and were hollowed out in a cup-shaped cavity in which the metal sample was placed.

The whole apparatus was connected up by a lead pipe to a set of trimount oil pumps in series, so that it could be quickly evacuated. The vacuum was tested by having a discharge tube sealed in by a side connection. The vacuum used when photographs were being taken was that which gives a dark, green discharge.

A hydrogen discharge tube fitted with a fluorite window was arranged in the apparatus, so that on one and the same plate there could be obtained both the hydrogen spectrum and the spectrum of any given metal in the same region. By adjusting the cover slit, a photograph of the gas spectrum could be obtained on the bottom of the plate, and then, without moving the plate, the cover slit could be readjusted, the discharge tube could be replaced by the vacuum arc lamp, and thus the spectrum of the metal could be thrown on the same plate just above the hydrogen spectrum. This afforded a means of comparing each metal spectrum with that of the gas.

¹ Lyman: Spectroscopy of the Extreme Ultra-Violet (Longmans).

² Handke: Inaug. Diss. Berlin., Aug. 1909.

³ Wolff: Ann. de. Phys. 42 p. 825, 1913.

⁴ Saunders: Astro phys. Jour. 43, p. 234, 1916.

It is important to notice that in obtaining the spectra of metals, the secondary gas spectrum is always present. Thus, by having the gas spectrum immediately above, and on the same plate, the lines due to the gas that might otherwise be attributed to the metal, could be easily picked out. When an intermittent arc, obtained by using a small current, was used, the time of exposure was longer and the secondary spectrum always came up strong, while when a high current was used, the arc remained much more steady and the secondary spectrum was much weaker relatively to that of the metal.

The length of the exposure ranged from 5 to 10 minutes for the metals. The steadier the arc remained, the shorter the time required. When being used, the arc could always be observed through a glass window sealed in the end of the casing of the arc.

The apparatus was connected up to a hydrogen tank, so that when not in use it could be filled with hydrogen gas. The hydrogen gas, together with phosphorous pentoxide kept the interior free from moisture.

RESULTS.

With the apparatus described above, the vacuum arc spectra of lead, tin, iron, nickel, cobalt, and thallium were investigated; also the spectra of copper, aluminium, zinc, and carbon, which were studied by Ainslie last year, were repeated, and the wave lengths of lines carefully measured again.

In calculating the wave lengths of the lines of these different spectra, certain lines previously determined were used as standards. From these lines, by means of graphical interpolation, the various lines were carefully measured. In working with a prism spectroscope it is necessary to use quite a large number of lines as standards, in order to get accurate results by graphical means. It is impossible to get results with this instrument by referring to a few standard lines, such as Saunders did using a grating spectroscope. For the region from 1850 down, hydrogen and aluminium lines measured by Lyman, and for the region above 1850, carbon monoxide by Lyman, and copper and thallium lines by Eder and Valenta, were used.

COPPER.

The copper vacuum arc spectrum was obtained by using a current of about 9 amperes. With this current the arc was almost continuous. The results obtained agree fairly well with those of Eder and Valenta for the region covered by their work and from λ 1750 down they agree with the values of Handke¹ obtained by using a copper spark.

¹ Lyman, Spectroscopy of the Extreme Ultra-Violet, p. 122.

| Intensity | Wave Length | $1/\lambda$ | E. & V. λ | Handke |
|-----------|-------------|-------------|----------------------|--------|
| 4 | 2370.0 | 42194 | 2369.94 | |
| 4 | 2297.0 | 43529 | | |
| 4 | 2267.0 | 44111 | | |
| 10 | 2247.0 | 44504 | 2247.08 | |
| 4 | 2231.5 | 44813 | | |
| 6 | 2221.0 | 45025 | | |
| 6 | 2213.0 | 45209 | | |
| 8 | 2195.8 | 45541 | 2195.87 | |
| 8 | 2182.2 | 45825 | 2181.80 | |
| | | | 2151.95 | |
| 4 | 2151.1 | 46488 | 2149.05 | |
| 6 | 2137.5 | 46784 | 2136.05 | |
| 4 | 2125.8 | 47041 | 2125.26 | |
| 4 | 2112.5 | 47337 | 2112.19 | |
| 4 | 2104.8 | 47510 | 2104.88 | |
| 4 | 2055.4 | 48652 | 2055.05 | |
| 6 | 2044.0 | 48724 | 2043.84 | |
| 6 | 2037.3 | 49084 | 2037.24 | |
| 4 | 2026.2 | 49353 | 2025.53 | |
| 4 | 2001.0 | 49975 | | |
| 4 | 1979.6 | 50515 | 1979.26 | 1749.9 |
| 4 | 1748.6 | 57189 | | 1747.1 |
| | | | | 1739.0 |
| 4 | 1739.7 | 57481 | | 1741.0 |
| 6 | 1721.8 | 58080 | | 1721.9 |
| 4 | 1708.5 | 58531 | | 1708.5 |
| 2 | 1704.9 | 58654 | | 1705.0 |
| 4 | 1693.4 | 59053 | | |
| 1 | 1692.5 | 59084 | | 1692.3 |
| 4 | 1686.7 | 59301 | | 1686.6 |
| 2 | 1684.6 | 59361 | | 1684.3 |
| | | | | 1681.2 |
| 1 | 1681.7 | 59428 | | 1681.9 |
| 2 | 1679.1 | 59556 | | 1679.0 |
| 2 | 1674.6 | 59680 | | 1674.5 |
| 2 | 1671.5 | 59826 | | 1671.6 |
| 2 | 1670.1 | 59877 | | 1669.8 |
| 2 | 1651.9 | 60536 | | 1651.9 |
| 8 | 1642.1 | 60898 | | 1641.8 |
| 8 | 1594.2 | 62727 | | 1594.2 |
| Intensity | .. | .. | E. & V. | Handke |

ALUMINIUM.

The vacuum arc spectrum for aluminium was obtained by using aluminium alone as terminals in the arc lamp. The arc remained steady for short periods but was mostly intermittent. For the intermittent arc the current was about eight amperes when the terminals touched together, but when separated the current quickly dropped to zero, and therefore it was necessary to separate and touch the terminals continually in order to obtain a photograph.

| Intensity | λ | $1/\lambda$ | λ |
|-----------|-----------|-------------|---------------|
| 10 | 2367.5 | 42239 | 2367.2 E & V. |
| 10 | 2139.5 | 46740 | |
| 7 | 2061.5 | 48508 | |
| 6 | 2026.0 | 49358 | |
| 14 | 1990.0 | 50251 | 1989.9 " |
| 4 | 1935.1 | 51677 | 1935.3 " |
| 10 | 1930.4 | 51803 | 1930.4 " |
| 32 | 1862.7 | 53685 | 1862.8 L |
| 32 | 1854.8 | 53914 | 1854.7 " |
| | | | {1766.9 H |
| 4 | 1766.6 | 56606 | {1766.0 " |
| 10 | 1762.0 | 56754 | 1761.9 L |
| 10 | 1724.3 | 57995 | 1725.0 L |
| 8 | 1720.7 | 58116 | 1721.2 " |
| 4 | 1718.5 | 58190 | 1719.3 " |
| 4 | 1670.6 | 59859 | 1670.6 " |
| 14 | 1611.7 | 62046 | 1611.8 " |
| 12 | 1605.6 | 62282 | 1605.6 " |

ZINC.

The vacuum arc spectrum of zinc was obtained by putting zinc in iron terminals in the arc lamp. The current used was 6 amperes which gave a fairly continuous and intense arc. The lines obtained do not agree with the values given by Handke¹ for a zinc spark.

| Int. | λ | $1/\lambda$ | λ |
|------|-----------|-------------|-----------------|
| 2 | 2372.5 | 42149 | |
| 2 | 2336.0 | 42808 | |
| 10 | 2287.5* | 43715 | |
| 6 | 2265.0* | 44150 | 2265.08 E. & V. |
| 2 | 2205.5 | 45341 | |
| 1 | 2171.5 | 46051 | |
| 16 | 2139.5 | 46739 | 2139.27 S |
| 2 | 2104.8 | 47510 | 2104.98 " |
| 4 | 2100.5 | 47608 | 2100.53 " |
| 2 | 2096.5 | 47698 | 2097.44 " |
| 6 | 2087.3 | 47908 | 2087.66 " |
| 1 | 2079.8 | 48082 | 2079.57 " |
| 2 | 2065.0 | 48426 | 2064.93 " |
| 6 | 2062.5 | 48485 | 2062.57 " |
| 16 | 2025.5 | 49370 | 2026.19 " |
| 3 | 1821.8 | 54890 | |
| 10 | 1589.6 | 62909 | 1589.76 W |
| 1 | 1510.4 | 66207 | |
| 1 | 1491.5 | 67047 | |
| 6 | 1486.2 | 67286 | 1486.20 W |
| 2 | 1478.5 | 67636 | |
| 4 | 1477.6 | 67923 | |
| 4 | 1457.9 | 68592 | 1457.9 W |
| 4 | 1457.5 | 68658 | 1457.56 " |
| 4 | 1451.1 | 69913 | 1450.82 " |
| 3 | 1445.0 | 69204 | |
| Int. | .. | .. | .. |

S. Saunders, Astrophysical Journal Vol. 43, p. 239, 1916. W. Wolf-Lyman, Spectroscopy of the Extreme Ultra-Violet, p. 123.

¹ These two lines may be due to cadmium but they appear to be zinc.

CARBON.

The carbon vacuum arc spectrum was obtained by using carbon alone as terminals in the vacuum arc lamp. A continuous arc was obtained by using a current of about 10 amperes. In addition to the lines given below a large number of broad and poorly defined bands occur which seem to correspond to those of the carbon monoxide spectrum. Upon measuring some of the more prominent ones, it was found that their wave lengths were different from those of the carbon monoxide spectrum obtained from the discharge tube. This may be due to the fact that the carbon arc is at a very high temperature compared to that of the discharge tube.

The lines of the spectra of copper, aluminium, zinc and carbon were measured last year by Ainslie, using the same apparatus but without the hydrogen comparison spectrum. The measurements were not very accurate and were repeated this year, using standard gas lines for comparison.

The carbon line spectrum is very strong in this region and the lines 1548.5, 1550.7 (1560.5, 1561.2, 1562.0) and 1656.9 correspond apparently to 1548.2, 1550.8, 1561.2 and 1656.8, given by Lyman as lines of certain origin. The line at 1561 appeared as a poorly defined triplet and was found in the spectra of all the metals worked with except zinc.

LINE SPECTRUM OF CARBON

| Int. | λ | $1/\lambda$ | λ |
|------|-----------|-------------|-----------------|
| 2 | 2307.5 | 43337 | 2296.94 E. & V. |
| 9 | 2298.0 | 43516 | |
| 2 | 2219.0 | 45065 | |
| 5 | 2088.5 | 47876 | 1930.12 " |
| 15 | 1930.5 | 51827 | |
| 9 | 1758.1 | 56879 | |
| 5 | 1749.7 | 57152 | |
| 10 | 1656.9 | 60354 | |
| 9 | 1562.0 | 64020 | |
| 9 | 1561.2 | 64053 | |
| 9 | 1560.5 | 64082 | |
| 2 | 1550.7 | 64487 | |
| 3 | 1548.5 | 64591 | |
| 5 | 1482.8 | 67439 | |
| 6 | 1464.5 | 68282 | |

BAND SPECTRUM OF CARBON

| Intensity | λ | Int. | λ |
|-----------|-----------|------|-----------|
| 3 | 2194.5 | 4 | 1811.7 |
| 4 | 2142.3 | 2 | 1806.8 |
| 4 | 2115.3 | 4 | 1793.6 |
| 4 | 2088.0 | 2 | 1774.6 |
| 3 | 2067.0 | 1 | 1748.0 |
| 2 | 2047.5 | 5 | 1729.9 |
| 2 | 2037.0 | 2 | 1725.0 |
| 1 | 2030.7 | 2 | 1721.3 |
| 4 | 1993.2 | 4 | 1712.7 |
| 3 | 1973.3 | 4 | 1706.5 |
| 2 | 1953.5 | 2 | 1670.6 |
| 2 | 1900.2 | 3 | 1653.4 |
| 1 | 1843.1 | 3 | 1649.1 |
| 1 | 1827.0 | 4 | 1630.7 |
| 2 | 1825.1 | 4 | 1597.9 |
| 1 | 1819.8 | 3 | 1576.6 |

IRON

The vacuum arc spectrum of iron was obtained by using iron terminals in the vacuum arc lamp. A current of 15 amperes was used, but even with this high current the arc was intermittent and not very bright, which made it difficult to get a good photograph of the spectrum.

| Int. | λ | $1/\lambda$ | λ |
|------|-----------|-------------|-----------------|
| 5 | 2394.5 | 41762 | 2394.68 E. & V. |
| 5 | 2380.2 | 42013 | |
| 4 | 2360.0 | 42373 | |
| 4 | 2346.0 | 42626 | 2345.48 " |
| 2 | 2097.5 | 47666 | 2097.20 " |
| 4 | 2078.8 | 48105 | |
| 3 | 2061.5 | 48508 | |
| 2 | 1926.0 | 51921 | 1925.60 B. |
| 4 | 1913.8 | 52252 | 1913.40 " |
| 3 | 1894.3 | 52790 | 1894.90 " |

B. Bloch, Journal de Physique, 5 Série, p. 628, 1914.

TIN.

To obtain the vacuum arc spectrum of tin, some pure tin metal was put in iron terminals in the vacuum arc lamp, the current was 5 amperes, which gave an intense intermittent arc. Lines were found in this spectrum. to extend far down in the ultra-violet region.

| Int. | λ | $1/\lambda$ | E. & V. λ | Saunders λ |
|------|-----------|-------------|----------------------|-----------------------|
| 6 | 2334.0 | 42845 | 2334.87 | 2335.53 |
| 6 | 2317.8 | 43144 | 2217.38 | 2317.93 |
| 3 | 2287.0 | 43725 | 2286.8 | 2287.28 |
| 8 | 2268.8 | 44075 | 2269.02 | 2269.65 |
| 6 | 2247.0 | 44504 | 2246.11 | 2246.73 |
| 6 | 2211.0 | 45228 | | 2210.38 |
| 5 | 2199.5 | 45465 | { 2199.68 | 2199.93 |
| | | | 2199.4 | |
| 12 | 2152.5 | 46457 | 2151.62 | 2152.08 |
| 2 | 2041.2 | 48990 | | |
| 1 | 1941.0 | 51520 | | |
| 20 | 1899.8 | 52637 | 1899.8 | |
| 6 | 1831.4 | 54603 | | 1811.29 |
| 20 | 1811.2 | 55212 | | |
| 16 | 1756.6 | 56928 | | |
| 1 | 1741.3 | 57428 | | |
| 20 | 1699.5 | 58841 | | |
| 6 | 1489.2 | 67150 | | |
| 9 | 1475.2 | 67787 | | |
| 4 | 1438.3 | 69526 | | |
| 4 | 1437.3 | 69574 | | |
| 4 | 1402.4 | 71303 | | |
| 4 | 1400.5 | 71402 | | |

THALLIUM.

The thallium vacuum arc spectrum was obtained by putting thallium in iron terminals. The current used was about 6 amperes which gave a continuous and intense arc. There is considerable variation in these results from those of Saunders. This may be due to the fact that Saunders used standard aluminium and zinc lines, while in this work the gas lines measured by Lyman were used as standards.

| Int. | λ | $1/\lambda$ | λ Saunders | λ E. & V. |
|------|-----------|-------------|-----------------------|----------------------|
| 4 | 2394.7 | 41759 | | 2394.72 |
| 4 | 2379.5 | 42043 | 2380.34 | 2379.68 |
| 2 | 2316.0 | 43178 | | 2316.14 |
| 8 | 2298.5 | 43506 | 2298.05 | 2298.25 |
| 2 | 2238.0 | 44683 | 2238.59 | |
| 2 | 2210.7 | 45235 | 2210.46 | |
| 2 | 2168.5 | 46115 | | |
| 2 | 2139.5 | 46740 | 2139.98 | 2139.44 |
| 14 | 1907.8 | 52416 | 1908.68 | |
| 6 | 1891.8 | 52860 | 1892.72 | |
| 2 | 1827.3 | 54725 | 1828.00 | |
| 6 | 1814.2 | 55121 | 1814.72 | |
| ■ | 1792.2 | 55797 | | |
| 6 | 1660.0 | 60241 | | |
| 6 | 1653.8 | 60467 | | |
| 8 | 1561.8 | 64029 | | |
| 16 | 1559.0 | 64144 | | |
| 5 | 1538.5 | 64998 | | |
| 4 | 1508.2 | 66304 | | |
| 6 | 1499.8 | 66676 | | |
| 2 | 1491.0 | 67069 | | |
| 4 | 1478.0 | 67659 | | |

LEAD.

The lead vacuum arc spectrum was obtained by putting lead in iron terminals in the vacuum arc lamp. The arc was almost continuous and very intense, being sustained by a current of 4 amperes. As it was not possible to obtain chemically pure lead, a sample of ordinary lead was used. Hence some of the lines recorded here may be due to impurities.

| Int. | λ | $1/\lambda$ | λ Saunders |
|------|-----------|-------------|-----------------------|
| 4 | 2430.5 | 41144 | |
| 5 | 2403.0 | 41515 | 2402.62 |
| 7 | 2394.0 | 41771 | 2394.52 |
| 5 | 2246.2 | 44520 | 2247.53 |
| 12 | 2204.4 | 45360 | 2204.18 |
| 8 | 2170.5 | 46072 | 2170.60 |
| 8 | 2060.5 | 48532 | |
| 2 | 1925.8 | 51926 | |
| 4 | 1913.7 | 52255 | |
| 2 | 1904.2 | 52527 | 1904.88 |
| 2 | 1898.7 | 52668 | |
| 2 | 1895.5 | 52756 | |
| 14 | 1821.7 | 54894 | 1822.06 |
| 10 | 1796.5 | 55664 | 1796.53 |
| 1 | 1744.2 | 57333 | |
| 2 | 1741.1 | 57435 | |
| 10 | 1726.2 | 57930 | |
| 12 | 1682.5 | 59435 | 1682.54 |
| 12 | 1671.6 | 59823 | |
| 3 | 1597.6 | 62594 | |
| 12 | 1555.8 | 64276 | |
| 4 | 1511.7 | 66150 | |
| 1 | 1494.7 | 66903 | |
| 1 | 1492.7 | 66993 | |
| 5 | 1434.0 | 69735 | |
| 5 | 1431.9 | 69837 | |

NICKEL.

The vacuum arc spectrum of nickel was obtained in a similar manner to that of iron. Nickel alone was put in the terminals of the vacuum arc lamp. The current used was 7 amperes, when the circuit was closed. This gave a strong intermittent arc. A large number of sharp and well-defined lines appear in the nickel spectrum in this region.

| Int. | λ | $1/\lambda$ | λ E. & V. |
|------|-----------|-------------|-------------------|
| 4 | 2394.0 | 41771 | 2394.68 |
| 4 | 2375.0 | 42105 | 2375.51 |
| 2 | 2366.5 | 42257 | 2366.62 |
| 2 | 2356.5 | 42436 | 2356.49 |
| 6 | 2345.5 | 42636 | 2345.48 |
| 6 | 2335.0 | 42827 | 2334.68 |
| 12 | 2315.5 | 43187 | 2316.12 |
| 12 | 2303.5 | 43412 | 2303.10 |
| 12 | 2297.5 | 43526 | 2297.60 |
| 7 | 2288.0 | 43706 | 2287.74 |
| 8 | 2278.8 | 43884 | 2278.65 |
| 6 | 2271.0 | 44033 | 2270.33 |
| 6 | 2264.5 | 44160 | 2264.57 |
| 8 | 2255.0 | 44346 | 2253.94 |
| 8 | 2226.4 | 44916 | 2226.41 |
| 8 | 2218.3 | 45080 | 2206.81 |
| 4 | 2208.2 | 45286 | 2206.72 |
| 4 | 2203.3 | 45386 | 2201.51 |
| 3 | 2195.2 | 45554 | |
| 6 | 2187.5 | 45714 | |
| 8 | 2177.7 | 45920 | |
| 9 | 2168.5 | 46115 | 2169.19 |
| 2 | 2161.0 | 46275 | 2161.31 |
| 5 | 2129.3 | 46964 | 2128.67 |
| 6 | 2114.3 | 47297 | 2113.61 |
| 6 | 2108.3 | 47432 | 2108.04 |
| 1 | 2097.7 | 47671 | 2097.2 |
| 2 | 2030.0 | 49261 | |
| 4 | 2021.0 | 49480 | |
| 1 | 2005.0 | 49875 | |
| 6 | 1991.0 | 50226 | |
| 2 | 1982.0 | 50454 | |
| 4 | 1964.8 | 50895 | |
| 2 | 1902.3 | 52568 | |
| 4 | 1859.2 | 53786 | |
| 6 | 1854.5 | 53923 | |
| 5 | 1849.7 | 54063 | |
| 6 | 1846.7 | 54151 | |

| Int. | λ | $1/\lambda$ | λ E.&V. |
|------|-----------|-------------|-----------------|
| 5 | 1829.4 | 54663 | |
| 4 | 1822.5 | 54870 | |
| 4 | 1818.7 | 54984 | |
| 1 | 1806.7 | 55350 | |
| 4 | 1794.3 | 55732 | |
| 4 | 1790.5 | 55850 | |
| 4 | 1787.5 | 55944 | |
| 2 | 1781.2 | 56126 | |
| 9 | 1767.8 | 56568 | |
| 5 | 1763.2 | 56715 | |
| 1 | 1758.7 | 56860 | |
| 4 | 1750.8 | 57117 | |
| 6 | 1746.0 | 57274 | |
| 4 | 1740.2 | 57465 | |
| 5 | 1737.3 | 57561 | |
| 7 | 1732.3 | 57727 | |
| 7 | 1729.4 | 57824 | |
| 3 | 1722.2 | 58065 | |
| 4 | 1720.0 | 58140 | |
| 5 | 1714.8 | 58282 | |
| 5 | 1709.5 | 58497 | |
| 3 | 1707.3 | 58572 | |
| 7 | 1701.3 | 58779 | |
| 2 | 1698.3 | 58882 | |
| 4 | 1692.5 | 59084 | |
| 4 | 1687.9 | 59245 | |
| 2 | 1661.8 | 60176 | |
| 2 | 1656.5 | 60368 | |
| 4 | 1653.2 | 60489 | |
| 7 | 1650.1 | 60569 | |

Bloch has also measured the spectrum of Nickel down to $\lambda 1851^1$. He has measured a great number of lines not obtained in this work. The strong lines of his spectrum correspond fairly well with the strong lines measured above.

COBALT.

The vacuum arc spectrum of Cobalt was obtained by putting cobalt in iron terminals. The current used was about 12 amperes when the circuit was closed. This gave an intense intermittent arc. The spectrum obtained was similar to that of iron and showed up lines that belonged to Carbon. Eder and Valenta's work covers the lines measured in this work down to $\lambda 2173.44$. Bloch has also made

¹Journal de Physique, 5 Série, p. 631, 1914.

measurements down to $\lambda 1872.94$, but his results do not correspond closely with those given below. This may be due to the fact that he used a spark in air, rather than the vacuum arc as used in this work.

| Intensity | λ | $1/\lambda$ | |
|-----------|-----------|-------------|--|
| ■ | 2387.5 | 41885 | |
| ■ | 2354.0 | 42481 | |
| ■ | 2347.0 | 42607 | |
| ■ | 2344.5 | 42653 | |
| 4 | 2325.5 | 43001 | |
| 7 | 2313.2 | 43230 | |
| 7 | 2308.0 | 43327 | |
| 2 | 2293.0 | 43611 | |
| 8 | 2286.5 | 43735 | |
| 4 | 2246.5 | 44534 | |
| 4 | 2138.7 | 46757 | |
| ■ | 2099.3 | 47635 | |
| 5 | 2061.5 | 48503 | |
| 7 | 2026.2 | 49354 | |
| 2 | 1939.5 | 51560 | |
| ■ | 1929.5 | 51827 | |
| 3 | 1912.2 | 52296 | |
| ■ | 1893.8 | 52804 | |
| 10 | 1861.4 | 53721 | |
| 10 | 1853.0 | 53966 | |
| ■ | 1819.8 | 54951 | |
| 1 | 1740.3 | 57461 | |
| ■ | 1710.9 | 58449 | |
| 7 | 1669.9 | 59384 | |

From $\lambda 2400$ to $\lambda 2000$ the experimental error is probably as much as one angstrom unit. This is due to the fact that the dispersion in this region is small, being about .05mm. per angstrom unit; also the lines obtained on the photographic plate are not as well defined. These measurements are given for a guide for future research work with vacuum arc sources.

From $\lambda 2000$ down, the probable error is on the average, well within half an angstrom unit. The dispersion varies from .067mm. at $\lambda 2000$ to .20mm. at $\lambda 1400$ per angstrom unit.

Summary of Results.

The vacuum arc spectra of copper, zinc, aluminium, carbon, iron, tin, lead, thallium, nickel and cobalt have been studied from $\lambda 2400$ to $\lambda 1400$. The vacuum arc spectra obtained for copper, zinc and aluminium were found to correspond with the results obtained

for the spark spectra of these metals by previous observers. For tin, lead and thallium, Saunders has given measurements between about $\lambda 2400$ and $\lambda 1800$, which correspond fairly well with these results. Several new lines have been observed and measured between $\lambda 1800$ and $\lambda 1400$. For the arc spectra of iron, cobalt, nickel and carbon, several lines were obtained, and measured, some of which corresponded with results given in previous work.

This work was begun by Mr. Ainslie under the direction of Professor J. C. McLennan and continued by the joint authors under the direction of Professor E. F. Burton.

UNIVERSITY OF TORONTO STUDIES

PAPERS FROM THE PHYSICAL LABORATORIES

The "Papers from the Physical Laboratories", issued as a special series of University of Toronto Studies, date from the year 1900. Nos. 1-17 were published by the Physical Department in a very limited edition and are no longer in print. For the sake of a complete record the numbering of the Papers, as forming a series of University of Toronto Studies, is made continuous with the earlier series and commences with No. 18. The earlier numbers, except those given below, are not now available either for sale or gift.

| | |
|---|------|
| No. 23: On the constitution and properties of Heusler's alloys, by H. A. McTAGGART | 0.50 |
| No. 24: Note on an improvement in the method of determining of visibility curves, by C. S. WRIGHT | 0.25 |
| No. 25: Note on the temperature variations in the specific re- sistance of Heusler's alloys, by H. A. McTAGGART and J. K. ROBERTSON | 0.25 |
| No. 26: On the conductivity of mixtures of dilute solutions, by J. A. GARDINER | 0.25 |
| No. 27: The absorption of the different types of Beta rays together with a study of the secondary rays excited by them, by V. E. POUND | 0.25 |
| No. 28: On variations on the conductivity of air enclosed in metallic receivers, by C. S. WRIGHT | 0.25 |
| No. 29: On charges gained by insulated metallic conductors, surrounded by other conductors, and the relation of these charges to the Volta effect, by J. K. ROBERTSON. | 0.25 |
| No. 30: On the radioactivity of potassium and other alkali metals, by Prof. J. C. McLENNAN and W. T. KENNEDY.. | 0.25 |
| No. 31: The action of electrolyte on copper colloidal solutions, by E. F. BURTON | 0.25 |
| No. 32: On the active deposit from actinium in uniform electric fields, by W. T. KENNEDY | 0.25 |
| No. 33: On the electric charges acquired in high vacua by insulated potassium salts and other radio-active sub- stances, by Professor J. C. McLENNAN | 0.25 |
| No. 34: On an improvement in the method of determining minimum spark potentials, by H. S. FIERHELLER | 0.25 |
| No. 35: On the penetrating radiation at the surface of the earth, by G. A. CLINE | 0.25 |
| No. 36: On the physical aspect of colloidal solution, by E. F. BURTON | 1.00 |
| No. 37: On a variation in the intensity of the penetrating radia- tion at the earth's surface observed during the passage of Halley's Comet, by ARTHUR THOMSON | 0.25 |
| No. 38: On the ionisation by collision in the gases helium and argon, by H. F. DAWES | 0.50 |
| No. 39: Effect of electricity on streams of water drops, by E. F. BURTON and W. B. WIEGAND | 0.25 |

| | | |
|---------|---|------|
| No. 40: | On the secondary rays excited by the alpha rays from polonium, Parts I and II, by V. E. POUND | 0.25 |
| No. 41: | The intensity of the earth's penetrating radiation at different altitudes and a secondary radiation excited by it, by Professor J. C. McLENNAN and E. N. MACALLUM | 0.25 |
| No. 42: | On the relation between the adiabatic and isothermal Young's moduli of metals, by E. F. BURTON | 0.25 |
| No. 43: | On the rotatory dispersion of quartz, by H. F. DAWES | 0.25 |
| No. 44: | On the mobilities of ions in gases at high pressures, by A. J. DEMPSTER | 0.25 |
| No. 45: | Measurements of precision on the penetrating radiation from the earth, by ARTHUR THOMSON | 0.25 |
| No. 46: | On the number of delta particles expelled concurrently with each alpha particle emitted by polonium, by W. T. KENNEDY | 0.25 |
| No. 47: | On the mobilities of ions in air at high pressures, by Professor J. C. McLENNAN and DAVID A. KEYS | 0.25 |
| No. 48: | On the absorption spectra of mercury, cadmium, zinc, and other metallic vapours, by Professor J. C. McLENNAN and EVAN EDWARDS | 0.50 |
| No. 49: | On the infra-red emission spectrum of the mercury arc, by Professor J. C. McLENNAN and RAYMOND C. DEARLE | 0.25 |
| No. 50: | On the ionisation tracks of alpha rays in hydrogen, by Professor J. C. McLENNAN and H. N. MERCER | 0.50 |
| No. 51: | On the delta rays emitted by zinc when bombarded by alpha rays, by Professor J. C. McLENNAN and C. G. FOUND | 0.25 |
| No. 52: | On the ultra-violet spectrum of elementary silicon by Professor J. C. McLENNAN and EVAN EDWARDS | 0.25 |
| No. 53: | On the ionisation potentials of magnesium and other metals, and on their absorption spectra, by Professor J. C. McLENNAN | 0.25 |
| No. 54: | On the Bunsen flame spectra of metallic vapours, by Professor J. C. McLENNAN and ANDREW THOMSON | 0.25 |
| No. 55: | On the ionisation of metallic vapours in flames, by Professor J. C. McLENNAN and DAVID A. KEYS | 0.25 |
| No. 56: | Emission and absorption in the infra-red spectrum of mercury, by RAYMOND C. DEARLE | 0.25 |
| No. 57: | Some experiments on residual ionization, by K. H. KINGDON | 0.25 |
| No. 58: | Image formation by Crystalline media, by H. F. DAWES | 0.25 |
| No. 59: | Periodic precipitation, by Miss A. W. FOSTER | 0.25 |
| No. 60: | Regularities in the spectra of lead and tin, by R. V. ZUMSTEIN | 0.25 |
| No. 61: | New lines in the extreme ultra-violet of certain metals, by D. S. AINSLIE and D. S. FULLER | 0.25 |



